

Computation of a Feynman integral and some identities of Clausen function values.

Gert Almkvist

Introduction.

For the 3-loop tetrahedral Feynman diagram with non-adjacent lines carrying masses a and b one gets the integral (see Broadhurst [1])

$$C(a, b) = -\frac{16}{b} \int_2^{2+b} \frac{dw}{w(w+a)\sqrt{w^2+b^2-4}} \arctan h \frac{w^2-4-2b}{w\sqrt{w^2+b^2-4}} \\ - \frac{16}{b} \int_{2+b}^{\infty} \frac{dw}{w(w+a)\sqrt{w^2+b^2-4}} \arctan h \frac{b}{\sqrt{w^2+b^2-4}}$$

Broadhurst also made the conjecture

$$C(1, 1) = 4\sqrt{2}(Cl_2(4\alpha) - Cl_2(2\alpha))$$

where $\sin(\alpha) = \frac{1}{3}$ and

$$Cl_2(x) = - \int_0^x \log(2 \left| \sin\left(\frac{\varphi}{2}\right) \right|) d\varphi = \sum_{n=1}^{\infty} \frac{\sin(nx)}{n^2}$$

is the Clausen function. I computed the two integrals (for $a = b = 1$) by hand

and found using the LLL-algorithm the following identities

Conjecture 1. Let $\tan(\alpha) = \frac{1}{\sqrt{2}}$ and $\tan(\beta) = \sqrt{8} + \sqrt{3}$ Then

$$1.1. Cl_2(\alpha) + Cl_2(\pi - \alpha) + Cl_2(\frac{\pi}{3} - \alpha) - Cl_2(\frac{2\pi}{3} - \alpha) = \frac{7}{4} Cl_2(\frac{2\pi}{3})$$

$$1.2. Cl_2(6\alpha - \pi) + Cl_2(\pi + 2\alpha) - 2Cl_2(2\alpha) + 2Cl_2(\pi - 4\alpha) = 0$$

$$1.3. Cl_2(\pi - 2\beta) + Cl_2(2\beta - 4\alpha) + Cl_2(2\beta - 2\alpha) - Cl_2(2\beta + 2\alpha - \pi) \\ - Cl_2(2\alpha) - 2Cl_2(\pi - 4\alpha) - 2Cl_2(\pi + 2\alpha) = 0$$

$$1.4. -12Cl_2(2\beta - 2\alpha) + 4Cl_2(\pi - 4\alpha) - 12Cl_2(\pi - 2\beta) - 18Cl_2(\pi + 2\alpha) + 7Cl_2(4\alpha) = 0$$

The first three identities give Broadhurst's value of $C(1, 1)$. But I could not prove any of the identities.

In July 1998 I met Broadhurst in Vancouver and he gave me [2] where he finds the value of $C(a, b)$ for general a and b and also proves it is correct. But I wanted to prove 1.1-1.4 and decided to compute $C(a, b)$ for $a^2 + b^2 < 4$ using elementary methods. The result contained 32 different Clausen values. Putting $a = \frac{1}{\pi}$ and $b = \frac{1}{e}$ and using the LLL-algorithm I found two general relations that specialized to 1.1 and 1.3 when $a = b = 1$. These two identities were easily proved by differentiation. 1.2 follows from an identity in [2] but 1.4 seems very difficult to prove (it is not a consequence of any general identity I found)

2. Computation of the integrals.

We start with two Lemmata.

Lemma 1. If $a^2 + b^2 \geq c^2$ and

$$\delta_1 = 2 \arctan\left(\frac{a+c}{b+\sqrt{a^2+b^2-c^2}}\right)$$

$$\delta_2 = 2 \arctan\left(\frac{a+c}{-b+\sqrt{a^2+b^2-c^2}}\right)$$

then

$$\begin{aligned} & \int_{\alpha}^{\beta} \log(a \cos(\varphi) + b \sin(\varphi) + c) d\varphi \\ &= (\beta - \alpha) \log\left(\frac{\sqrt{a^2+b^2}}{2}\right) + Cl_2(\delta_2 - \beta) - Cl_2(\delta_2 - \alpha) + Cl_2(\delta_1 + \alpha) - Cl_2(\delta_1 + \beta) \end{aligned}$$

Proof: We have

$$a \cos(\varphi) + b \sin(\varphi) + c = 2\sqrt{a^2+b^2} \sin\left(\frac{\delta_2 - \varphi}{2}\right) \sin\left(\frac{\delta_1 + \varphi}{2}\right)$$

Taking logarithms and integrating we are done.

Lemma 2:

$$\begin{aligned} & \int_{\alpha}^{\beta} \log(\tan(\varphi) - \tan(\delta)) d\varphi \\ &= \frac{1}{2} \{Cl_2(2\alpha - 2\delta) - Cl_2(2\beta - 2\delta) + Cl_2(\pi - 2\alpha) - Cl_2(\pi - 2\beta)\} - (\beta - \alpha) \log(\cos(\delta)) \end{aligned}$$

Proof: We have

$$\log(\tan(\varphi) - \tan(\delta)) = \log(2 \sin(\varphi - \delta)) - \log(2 \cos(\varphi)) - \log(\cos(\delta))$$

Then use

$$\int_{\alpha}^{\beta} \log(2 \cos(\varphi)) d\varphi = \frac{1}{2} \{Cl_2(\pi - 2\beta) - Cl_2(\pi - 2\alpha)\}$$

Using the partial fraction

$$\frac{1}{w(w+a)} = \frac{1}{a} \left(\frac{1}{w} - \frac{1}{w+a} \right)$$

we get two integrals not containing a

$$I_1 = \int_{2+b}^{\infty} \frac{dw}{w\sqrt{w^2+b^2-4}} \arctan h\left(\frac{b}{\sqrt{w^2+b^2-4}}\right)$$

$$I_2 = \int_2^{2+b} \frac{dw}{w\sqrt{w^2+b^2-4}} \arctan h\left(\frac{w^2-2(2+b)}{w\sqrt{w^2+b^2-4}}\right)$$

Put $c = \sqrt{4-b^2}$ and make the substitution

$$w = \frac{c}{\sin(\varphi)}$$

and use the notation

$$\begin{aligned} \sin(\alpha_1) &= \sqrt{\frac{2-b}{2+b}} \\ \tan(\alpha_2) &= \frac{c}{b} \end{aligned}$$

Then

$$\begin{aligned} I_1 &= \frac{1}{2c} \int_0^{\alpha_1} \log\left(\frac{c \cos(\varphi) + b \sin(\varphi)}{c \cos(\varphi) - b \sin(\varphi)}\right) d\varphi \\ &= \frac{1}{4c} \{-Cl_2(2\alpha_1 + 2\alpha_2) + Cl_2(2\alpha_1 - 2\alpha_2) + 2Cl_2(2\alpha_2)\} \\ &= \frac{1}{4c} \{-q_1 + q_2 + 2q_3\} \end{aligned}$$

Similarly using Lemma 1 and 2

$$\begin{aligned} I_2 &= \frac{1}{2c} \int_{\alpha_1}^{\alpha_2} \left\{ \log\left(\frac{\cos(\varphi) - \frac{b}{2}}{\cos(\varphi) + \frac{b}{2}}\right) - 2 \log\left(\tan\left(\frac{\varphi}{2}\right)\right) \right\} \\ &= \frac{1}{2c} \left\{ \begin{array}{l} -Cl_2(\alpha_2 - \alpha_1) + Cl_2(\alpha_2 + \alpha_1) - Cl_2(2\alpha_2) - Cl_2(\pi - 2\alpha_2) + Cl_2(\pi - \alpha_1 - \alpha_2) \\ -Cl_2(\pi + \alpha_1 - \alpha_2) + 2Cl_2(\alpha_2) - 2Cl_2(\alpha_1) + 2Cl_2(\pi - \alpha_2) - 2Cl_2(\pi - \alpha_1) \end{array} \right\} \\ &= \frac{1}{2c} \{-q_4 + q_5 - q_6 - q_7 + q_8 - q_9 + 2q_{10} - 2q_{11} + 2q_{12} - 2q_{13}\} \end{aligned}$$

We will show that

$$I_1 + I_2 = 0$$

Using the duplication formula

$$Cl_2(2x) = 2Cl_2(x) - 2Cl_2(\pi - x)$$

we find

$$q_1 = 2q_5 - 2q_8$$

$$q_2 = 2q_9 - 2q_4$$

and also

$$q_3 = q_6$$

It follows that $I_1 + I_2 = 0$ is equivalent to

$$2q_4 + q_7 - 2q_8 - 2q_{10} + 2q_{11} - 2q_{12} + 2q_{13} = 0$$

which follows from the following result ($\alpha = \alpha_1, \beta = \alpha_2$)

Theorem 1: Assume $\sin(\alpha) = \tan(\frac{\beta}{2})$ Then

$$Cl_2(\pi - 2\beta) - 2Cl_2(\beta) - 2Cl_2(\pi - \beta) + 2Cl_2(\alpha) + 2Cl_2(\pi - \alpha) + 2Cl_2(\beta - \alpha) - 2Cl_2(\pi - \alpha - \beta) = 0$$

Proof: Since $Cl_2(\pi) = 0$ we find that the identity is true when $\alpha = \beta = 0$.
Let $\tan(\frac{\alpha}{2}) = t$ and consider α and β as functions of t . We have

$$\begin{aligned}\frac{d}{d\alpha}Cl_2(\alpha) &= -\log(2\sin(\frac{\alpha}{2})) \\ \frac{d}{d\alpha}Cl_2(\pi - \alpha) &= \log(\cos(\frac{\alpha}{2}))\end{aligned}$$

Let $LHS = f(t)$. Then

$$\begin{aligned}-2\frac{df}{dt} &= \left\{ -2\log(2\cos(\beta)) - 2\log(2\sin(\frac{\beta}{2})) + 2\log(2\cos(\frac{\beta}{2})) \right\} \beta' \\ &+ \left\{ 2\log(2\sin(\frac{\alpha}{2})) - 2\log(2\cos(\frac{\alpha}{2})) \right\} \alpha' + 2\log(2\sin(\frac{\beta - \alpha}{2}))(\beta' - \alpha') \\ &\quad + 2\log(2\cos(\frac{\beta + \alpha}{2}))(\beta' + \alpha') \\ &= \log\left(\frac{2\cos(\frac{\beta}{2})\sin(\frac{\beta - \alpha}{2})\cos(\frac{\beta + \alpha}{2})}{\sin(\frac{\beta}{2})\cos(\beta)}\right)\beta' + \log\left(\frac{\sin(\frac{\alpha}{2})\cos(\frac{\beta + \alpha}{2})}{\cos(\frac{\alpha}{2})\sin(\frac{\beta - \alpha}{2})}\right)\alpha'\end{aligned}$$

Using $\tan(\frac{\alpha}{2}) = t$ we obtain

$$\begin{aligned}\cos(\alpha) &= \frac{1 - t^2}{1 + t^2} \\ \sin(\alpha) &= \tan(\frac{\beta}{2}) = \frac{2t}{1 + t^2} \\ \cos(\beta) &= \frac{(1 - t^2)^2}{1 + 6t^2 + t^4} \\ \sin(\beta) &= \frac{4t(1 + t^2)}{1 + 6t^2 + t^4}\end{aligned}$$

Then

$$\begin{aligned}\frac{2\cos(\frac{\beta}{2})\sin(\frac{\beta - \alpha}{2})\cos(\frac{\beta + \alpha}{2})}{\sin(\frac{\beta}{2})\cos(\beta)} &= \frac{\sin(\beta) - \sin(\alpha)}{\tan(\frac{\beta}{2})\cos(\beta)} \\ &= \frac{\frac{4t(1 + t^2)}{1 + 6t^2 + t^4} - \frac{2t}{1 + t^2}}{\frac{2t}{1 + t^2} \cdot \frac{(1 - t^2)^2}{1 + 6t^2 + t^4}} = 1\end{aligned}$$

Similarly

$$\frac{\sin(\frac{\alpha}{2}) \cos(\frac{\beta+\alpha}{2})}{\cos(\frac{\alpha}{2}) \sin(\frac{\beta-\alpha}{2})} = 1$$

Hence $\frac{df}{dt} = 0$ and since $f(0) = 0$ we have $f(t) \equiv 0$

Corollarium (Conjecture 1.1): Assume $\sin(\alpha) = \frac{1}{\sqrt{3}}$. Then

$$Cl_2(\alpha) + Cl_2(\pi - \alpha) + Cl_2(\frac{\pi}{3} - \alpha) - Cl_2(\frac{2\pi}{3} - \alpha) = \frac{7}{4} Cl_2(\frac{2\pi}{3})$$

Proof: We get

$$\tan(\frac{\beta}{2}) = \sin(\alpha) = \frac{1}{\sqrt{3}}$$

and hence $\beta = \frac{\pi}{3}$. Using $Cl_2(\frac{\pi}{3}) = \frac{3}{2} Cl_2(\frac{2\pi}{3})$ and the Theorem we get the wanted result.

Since $I_1 + I_2 = 0$ we get

$$C(a, b) = \frac{16}{ab} \{I_3 + I_4\}$$

where

$$I_3 = \int_{2+b}^{\infty} \frac{dw}{(w+a)\sqrt{w^2-c^2}} \arctan h\left(\frac{b}{\sqrt{w^2-c^2}}\right)$$

$$I_4 = \int_2^{2+b} \frac{dw}{(w+a)\sqrt{w^2-c^2}} \arctan h\left(\frac{w^2-4-2b}{w\sqrt{w^2-c^2}}\right)$$

In I_3 we substitute

$$w+a = \frac{1}{u}$$

Let

$$p = a+b+2$$

$$d = \sqrt{4-a^2-b^2}$$

Then we get

$$I_3 = \int_0^{1/p} \frac{du}{\sqrt{1-2au-d^2u^2}} \arctan h\left(\frac{bu}{\sqrt{1-2au+d^2u^2}}\right)$$

$$= \frac{1}{2d} \int_{\alpha_3}^{\alpha_4} \log\left(\frac{dc \cos(\varphi) + bc \sin(\varphi) - ab}{dc \cos(\varphi) - bc \sin(\varphi) + ab}\right) d\varphi$$

after the substitution

$$u = \frac{c \sin(\varphi) - a}{d^2}$$

where

$$\sin(\alpha_3) = \frac{a}{c}$$

$$\sin(\alpha_4) = \frac{a}{c} + \frac{d^2}{cp}$$

Let

$$\delta_1 = 2 \arctan\left(\frac{cd - ab}{2d + bc}\right)$$

$$\delta_2 = 2 \arctan\left(\frac{cd - ab}{2d - bc}\right)$$

$$\delta_3 = 2 \arctan\left(\frac{cd + ab}{2d - bc}\right)$$

$$\delta_4 = 2 \arctan\left(\frac{cd + ab}{2d + bc}\right)$$

Using Lemma 2 we obtain

$$\begin{aligned} I_3 &= \frac{1}{2d} \left\{ \begin{array}{l} Cl_2(\delta_2 - \alpha_4) - Cl_2(\delta_2 - \alpha_3) + Cl_2(\delta_1 + \alpha_3) - Cl_2(\delta_1 + \alpha_4) \\ - Cl_2(\delta_4 - \alpha_4) + Cl_2(\delta_4 - \alpha_3) - Cl_2(\delta_3 + \alpha_3) + Cl_2(\delta_3 + \alpha_4) \end{array} \right\} \\ &= \frac{1}{2d} \{r_1 - r_2 + r_3 - r_4 - r_5 + r_6 - r_7 + r_8\} \end{aligned}$$

Finally we substitute

$$cu = w + \sqrt{w^2 - c^2}$$

to get

$$I_4 = \frac{1}{c} \int_{u_1}^{u_2} \frac{du}{u^2 + \frac{2a}{c}u + 1} \log\left(\frac{u^2 - f^2}{f^2u^2 - 1}\right)$$

where

$$f^2 = \frac{2 + b}{2 - b}$$

$$u_1 = f$$

$$u_2 = f + \sqrt{\frac{2b}{2 - b}}$$

Then put

$$u = \frac{d \tan(\varphi) - a}{c}$$

to get

$$I_4 = \frac{1}{d} \int_{\alpha_6}^{\alpha_7} \log\left(\frac{d^2(\tan(\varphi) - \frac{a}{d})^2(\tan(\varphi) - \frac{a+b+2}{d})(\tan(\varphi) - \frac{a-b-2}{d})}{c^2 f^2(\tan(\varphi) - \frac{a-b+2}{d})(\tan(\varphi) - \frac{a+b-2}{d})}\right) d\varphi$$

where

$$\tan(\alpha_6) = \frac{p}{d}$$

$$\tan(\alpha_7) = \frac{p + \sqrt{2b^2 + 4b}}{d}$$

Define

$$\delta_7 = \arctan\left(\frac{a}{d}\right)$$

$$\delta_8 = \arctan\left(\frac{a+b+2}{d}\right) = \alpha_6$$

$$\delta_9 = \arctan\left(\frac{a-b-2}{d}\right)$$

$$\delta_{10} = \arctan\left(\frac{a-b+2}{d}\right)$$

$$\delta_{11} = \arctan\left(\frac{a+b-2}{d}\right)$$

Using Lemma 2 we get (all logarithms cancel)

$$I_4 = \frac{1}{2d} \left\{ \begin{array}{l} 2Cl_2(2\alpha_6 - 2\delta_7) - 2Cl_2(2\alpha_7 - 2\delta_7) - Cl_2(2\alpha_7 - 2\delta_8) + Cl_2(2\alpha_6 - 2\delta_9) \\ -Cl_2(2\alpha_7 - 2\delta_9) - Cl_2(2\alpha_6 - 2\delta_{10}) + Cl_2(2\alpha_6 - 2\delta_{11}) + Cl_2(2\alpha_7 - 2\alpha_{11}) \\ + 2Cl_2(\pi - 2\alpha_6) - 2Cl_2(\pi - 2\alpha_7) \end{array} \right\}$$

$$= \frac{1}{2d} \{2r_9 - 2r_{10} - r_{11} + r_{12} - r_{13} - r_{14} + r_{15} - r_{16} + r_{17} + 2r_{18} - 2r_{19}\}$$

Using LLL on $\{r_1, r_2, \dots, r_{19}\}$ with the special values $a = \frac{1}{\pi}$ and $b = \frac{1}{e}$ we found

$$r_2 = r_9$$

$$r_5 = r_{11}$$

$$r_4 = -r_{13}$$

$$r_1 = r_{15}$$

$$r_8 = -r_{17}$$

$$r_6 = r_{18}$$

These identities are valid for all a and b such that $a^2 + b^2 < 4$. They are not trivial. E.g. $r_5 = r_{11}$ says that

$$\frac{1}{2}(\delta_4 - \alpha_4) = \alpha_7 - \delta_8$$

which is equivalent to (after taking $\tan$ on both sides)

$$\frac{du}{p^2 + d^2 + pu} = \frac{\frac{ab+cd}{2d+bc} - \frac{pc-du}{d^2+ap}}{1 + \frac{(ab+cd)(pc-du)}{(2d+bc)(d^2+ap)}}$$

where

$$\begin{aligned}c &= \sqrt{4 - b^2} \\d &= \sqrt{4 - a^2 - b^2} \\p &= a + b + 2 \\u &= \sqrt{2b^2 + 4b}\end{aligned}$$

Maple verifies this identity and the other relations as well. Using these identities we reduce $I_3 + I_4$ to eleven Clausen values. In order to simplify this and to get the same answer as Broadhurst got in [2] we have to introduce some more notation. Let

$$\begin{aligned}\phi &= \arctan\left(\frac{d}{p}\right) \\\phi_a &= \arctan\left(\frac{d}{a}\right) \\\phi_b &= \arctan\left(\frac{d}{b}\right) \\s_1 &= Cl_2(4\phi) \\s_2 &= Cl_2(2\phi_a + 2\phi_b - 2\phi) \\s_3 &= Cl_2(2\phi_a - 2\phi) \\s_4 &= Cl_2(2\phi_b - 2\phi) \\s_5 &= Cl_2(2\phi_a + 2\phi_b - 4\phi) \\s_6 &= Cl_2(2\phi_a) \\s_7 &= Cl_2(2\phi_b) \\s_8 &= Cl_2(2\phi)\end{aligned}$$

Then Broadhurst's result is

$$2d(I_3 + I_4) = s_1 + s_2 + s_3 + s_4 - s_5 - s_6 - s_7 - s_8$$

Comparing with our value for $I_3 + I_4$ this is equivalent to the following identities found by LLL

$$-2r_{10} - 2r_{11} - r_{14} + 2r_{15} - 2r_{19} - s_1 + s_6 + 4s_8 = 0$$

$$r_3 = s_4$$

$$r_7 = -s_2$$

$$r_9 = s_3$$

$$r_{12} = -s_7$$

$$r_{16} = s_5$$

$$r_{18} = s_8$$

The last six of these follow from

$$\alpha_3 = \frac{\pi}{2} - \phi_a$$

$$\alpha_6 = \frac{\pi}{2} - \phi$$

$$\delta_1 = -2\phi + \phi_a + 2\phi_b - \frac{\pi}{2}$$

$$\delta_3 = 2\phi - \phi_a - 2\phi_b + \frac{3\pi}{2}$$

$$\delta_7 = \frac{\pi}{2} - \phi_a$$

$$\delta_9 = -\phi + \phi_b - \frac{\pi}{2}$$

$$\delta_{11} = \frac{\pi}{2} + \phi - \phi_a - \phi_b$$

which are easily verified by Maple. It remains to prove the first identity. It follows from the following result.

Proposition 1: Let

$$d = \sqrt{4 - a^2 - b^2}$$

$$p = a + b + 2$$

$$\gamma = \arctan\left(\frac{p + \sqrt{2b^2 + 4b}}{d}\right)$$

$$\phi = \arctan\left(\frac{d}{p}\right)$$

$$\phi_a = \arctan\left(\frac{d}{a}\right)$$

Then

$$\begin{aligned} & 2Cl_2(2\gamma + 2\phi_a - \pi) + 2Cl_2(2\gamma + 2\phi - \pi) + Cl_2(2\phi_a - 4\phi) - 2Cl_2(2\gamma - 2\phi + 2\phi_a - \pi) \\ & + 2Cl_2(\pi - 2\gamma) + Cl_2(4\phi) - Cl_2(2\phi_a) - 4Cl_2(2\phi) = 0 \end{aligned}$$

Proof: consider the left hand side as a function $f(a, b)$. Differentiating we get

$$\begin{aligned} -2df &= 4\log(2\sin(\gamma + \phi_a - \frac{\pi}{2}))(d\gamma + d\phi_a) + 4\log(2\sin(\gamma + \phi - \frac{\pi}{2}))(d\gamma + d\phi) \\ & 2\log(2\sin(\phi_a - 2\phi))(d\phi_a - 2d\phi) - 4\log(2\sin(\gamma - \phi + \phi_a - \frac{\pi}{2}))(d\gamma - d\phi + d\phi_a) \\ & -4\log(2\sin(\frac{\pi}{2} - \gamma))d\gamma + 4\log(2\sin(2\phi))d\phi - 2\log(2\sin(\phi_a))d\phi_a - 8\log(2\sin(\phi))d\phi \\ & = 4\log\left(\frac{\sin(\gamma + \phi_a - \frac{\pi}{2})\sin(\gamma + \phi - \frac{\pi}{2})}{\sin^2(\gamma - \phi + \phi_a - \frac{\pi}{2})\sin(\phi_a)}\right)d\gamma \end{aligned}$$

$$\begin{aligned}
& +2 \log\left(\frac{\sin^2(\gamma + \phi_a - \frac{\pi}{2}) \sin(\phi_a - 2\phi)}{\sin^2(\gamma - \phi + \phi_a - \frac{\pi}{2}) \sin(\phi_a)}\right) d\phi_a \\
& +4 \log\left(\frac{\sin(\gamma + \phi - \frac{\pi}{2}) \sin(\gamma - \phi + \phi_a - \frac{\pi}{2}) \sin(2\phi)}{\sin(\phi_a - 2\phi) \sin^2(\phi)}\right) d\phi \\
& = 4 \log(T_1) d\gamma + 2 \log(T_2) d\phi_a + 4 \log(T_3) d\phi = 0
\end{aligned}$$

since $T_1 = T_2 = T_3 = 1$. We show that $T_3 = 1$ which is equivalent to

$$\cos(\gamma + \phi) \cos(\gamma + \phi_a - \phi) \sin(2\phi) = \sin(\phi_a - 2\phi) \sin^2(\phi)$$

if and only if

$$\begin{aligned}
& 2(1 - \tan(\gamma) \tan(\phi))(1 - \tan(\gamma) \tan(\phi_a) + \tan(\phi) \tan(\gamma) + \tan(\phi) \tan(\phi_a)) \\
& = \tan(\phi)(1 + \tan^2(\gamma))(\tan(\phi_a)(1 - \tan^2(\phi)) - 2 \tan(\phi))
\end{aligned}$$

if and only if

$$\begin{aligned}
& 2p\sqrt{2b^2 + 4b}(p^2 - d^2 - 2ap + (b + 2)\sqrt{2b^2 + 4b}) \\
& = (p^2 - d^2 - 2ap)(d^2 + p^2 + 2b^2 + 4b + 2p\sqrt{2b^2 + 4b})
\end{aligned}$$

which is easily verified. Similarly one verifies that $T_1 = T_2 = 1$. Hence $df = 0$ which means that $f(a, b)$ is constant. Putting $a = b = 0$ we get $d = 2, p = 2, \gamma = \phi = \frac{\pi}{4}, \phi_a = \frac{\pi}{2}$ which gives $f(0, 0) = 0$. It follows that $f(a, b) \equiv 0$ and the proof is finished.

Corollary: We have if $a^2 + b^2 < 4$

$$C(a, b) = \frac{8}{ab\sqrt{4 - a^2 - b^2}} \left\{ \begin{array}{l} Cl_2(4\phi) + Cl_2(2\phi_a + 2\phi_b - 2\phi) + Cl_2(2\phi_a - 2\phi) + Cl_2(2\phi_b - 2\phi) \\ -Cl_2(2\phi_a + 2\phi_b - 4\phi) - Cl_2(2\phi_a) - Cl_2(2\phi_b) - Cl_2(2\phi) \end{array} \right\}$$

Proposition 2: We have

$$\begin{aligned}
& 2Cl_2(2\phi) - 4Cl_2(2\phi_b) + Cl_2(4\phi_b) + 2Cl_2(2\phi_b - 2\phi) - 2Cl_2(2\phi_a - 2\phi) \\
& + Cl_2(2\phi_a - 4\phi) + 2Cl_2(2\phi_a + 2\phi_b - 2\phi) - Cl_2(2\phi_a + 4\phi_b - 4\phi) = 0
\end{aligned}$$

Proof: Let $f(a, b)$ be the LHS. Differentiating we get

$$\begin{aligned}
df & = 4 \log\left(\frac{\sin(\phi) \sin(\phi_a - \phi) \sin(\phi_a + 2\phi_b - 2\phi)}{\sin(\phi_b - \phi) \sin(\phi_a - 2\phi) \sin(\phi_a + \phi_b - \phi)}\right) d\phi \\
& + 2 \log\left(\frac{\sin(\phi_a - 2\phi) \sin^2(\phi_a + \phi_b - \phi)}{\sin^2(\phi_b - \phi) \sin(\phi_a + 2\phi_b - 2\phi)}\right) d\phi_a \\
& + 4 \log\left(\frac{\sin(2\phi_b) \sin(\phi_b - \phi) \sin(\phi_a + \phi_b - \phi)}{\sin^2(\phi_b) \sin(\phi_a + 2\phi_b - 2\phi)}\right) d\phi_b = 0
\end{aligned}$$

since all logarithms are zero (checked by Maple). Since $f(0, 0) = 0$ we have $f(a, b) \equiv 0$.

Corollarium 1: Identity 1.2 is true.

Proof: Put $a = b = 1$ in Proposition 2. Let $\alpha = \arctan(\frac{1}{\sqrt{2}})$. Then $\phi = \frac{\pi}{2} - 2\alpha$ and $\phi_a = \phi_b = \frac{\pi}{2} - \alpha$. Inserting this into Proposition 2 we obtain

$$2CL_2(\pi - 4\alpha) - 4Cl_2(\pi - 2\alpha) - Cl_2(4\alpha) + Cl_2(6\alpha + \pi) - Cl_2(\pi + 2\alpha) = 0$$

Using the duplication formula

$$Cl_2(4\alpha) = 2Cl_2(2\alpha) - 2Cl_2(\pi - 2\alpha)$$

we get 1.2

Corollarium 2. Identity 1.3 is true

Proof: With $a = b = 1$ in Proposition 1 we have $\tan(\gamma) = \sqrt{8} + \sqrt{3}$. Hence $\gamma = \beta$ in 1.3. Proposition 1 gives

$$2Cl_2(2\beta - 2\alpha) + 2Cl_2(2\beta - 4\alpha) + Cl_2(6\alpha - \pi) - 2Cl_2(2\beta + 2\alpha - \pi) + 2Cl_2(\pi - 2\beta) \\ + Cl_2(2\pi - 8\alpha) - Cl_2(\pi - 2\alpha) - 4Cl_2(\pi - 4\alpha) = 0$$

Using 1.2 for $Cl_2(6\alpha - \pi)$ and the duplication formula for $Cl_2(8\alpha)$ we obtain 1.3

Acknowledgements. The beginning of this work was done in 1998 at CECM, Simon Fraser University. I want to thank Jonathan and Peter Borwein for many stimulating discussions. The final part of the paper could not have been written without Broadhurst's result in [2]-

References:

1. D.Broadhurst, A dilogarithmic 3-dimensional Ising tetrahedron, hep-th/9805025
2. D.Broadhurst, Solving differential equations for 3-loop diagrams: relation to hyperbolic geometry and knot theory, hep-th/9806174.

Appendix. Proof of an identity found by Broadhurst

Let $\sin(\alpha) = \frac{1}{3}$. Then Broadhurst in [1] found the following identity using PSLQ

$$4\sqrt{2}(Cl_2(4\alpha) - Cl_2(2\alpha)) \\ = \sum_{n=0}^{\infty} \left(-\frac{1}{2}\right)^{3n} \frac{1}{n + \frac{1}{2}} \left(\frac{1}{n + \frac{1}{2}} - 3\log(2)\right) - 3 \sum_{n=1}^{\infty} \left(-\frac{1}{2}\right)^{3n} \frac{H_n}{n + \frac{1}{2}}$$

where

$$H_n = \sum_{k=1}^n \frac{1}{k}$$

We will prove this here using several formulas for the dilogarithm

$$Li_2(x) = \sum_{n=1}^{\infty} \frac{x^n}{n^2}$$

We start by quoting some formulas in Lewin [2] pp.244

Lemma 1

1.1.

$$Li_2(x) + Li_2(-x) = \frac{1}{2}Li_2(x^2)$$

1.2.

$$Li_2(x) + Li_2(-\frac{x}{1-x}) = -\frac{1}{2}\log^2(1-x)$$

1.3.

$$Li_2(\frac{1}{1+x}) - Li_2(-x) = \frac{\pi^2}{6} - \frac{1}{2}\log(1+x)\log(\frac{1+x}{x^2})$$

1.4.(Abel)

$$Li_2(\frac{x}{1-x} \cdot \frac{y}{1-y}) = Li_2(\frac{x}{1-y}) + Li_2(\frac{y}{1-x}) - Li_2(x) - Li_2(y) - \log(1-x)\log(1-y)$$

1.5.

$$Li_2(x) + Li_2(1-x) = \frac{\pi^2}{6} - \log(x)\log(1-x)$$

Lemma 2. We have

$$\begin{aligned} & \sum_{n=1}^{\infty} \frac{H_n}{2n+1} z^{2n+1} \\ &= \frac{1}{2} \left\{ \frac{1}{2} \log^2(1-z) - \frac{1}{2} \log^2(1+z) + \log(2) \log\left(\frac{1-z}{1+z}\right) + Li_2\left(\frac{1+z}{2}\right) - Li_2\left(\frac{1-z}{2}\right) \right\} \end{aligned}$$

Proof: We have by [3], p.715

$$\sum_{n=1}^{\infty} H_n x^{2n} = -\frac{\log(1-x^2)}{1-x^2}$$

Integrate

$$\sum_{N01}^{\infty} \frac{H_n}{2n+1} x^{2n+1} = -\frac{1}{2} \int_0^x (\log(1-t) + \log(1+t)) \left(\frac{1}{1-t} + \frac{1}{1+t} \right) dt$$

and we are done by formula 3.2 p.266 in Lewin [2]

$$\begin{aligned} & \int_0^x \frac{\log(a+bt)}{c+et} dt \\ &= \frac{1}{e} \left\{ \log\left(\frac{ae-bc}{e}\right) \log\left(\frac{c+ex}{c}\right) - Li_2\left(\frac{b(c+ex)}{bc-ae}\right) + Li_2\left(\frac{bc}{bc-ae}\right) \right\} \end{aligned}$$

Proof of Broadhurst's identity.

Introduce the notation

$$x = \frac{1}{2} \left(1 + \frac{i}{\sqrt{8}} \right)$$

$$y = \frac{1}{2}$$

$$u = \frac{\sqrt{8} + i}{3}$$

$$z = -\frac{i}{8}$$

Then we want to compute

$$Cl_2(4\alpha) - Cl_2(2\alpha) = \text{Im}(Li_2(u^4) - Li_2(u^2))$$

In Abel's identity (Lemma 1.4) we have

$$\frac{x}{1-x} = u^2, \quad \frac{y}{1-y} = 1, \quad \frac{x}{1-y} = 1-z, \quad \frac{y}{1-x} = \frac{1}{1+z}$$

and hence

$$(2.1) \quad Li_2(u^2) = Li_2(1-z) + Li_2\left(\frac{1}{1+z}\right) - Li_2(x) - Li_2\left(\frac{1}{2}\right) + \log(2) \log(1-x)$$

By Lemma 1.5 and 1.3 we get

$$(2.2) \quad Li_2(1-z) = -Li_2(z) + \frac{\pi^2}{6} - \frac{1}{2} \log(z) \log(1-z)$$

$$(2.3) \quad Li_2\left(\frac{1}{1+z}\right) = Li_2(-z) + \frac{\pi^2}{6} - \frac{1}{2} \log(1+z) \log\left(\frac{1+z}{z^2}\right)$$

Adding (2.1), (2.2) and (2.3) we obtain

$$(2.4) \quad Li_2(u^2) = Li_2(\bar{z}) - Li_2(z) - Li_2(x) + \frac{\pi^2}{3} - Li_2\left(\frac{1}{2}\right) + \log(2) \log(1-x) - \log(z) \log(1-z) - \frac{1}{2} \log(1+z) \log\left(\frac{1+z}{z^2}\right)$$

By Lemma 1.2 and 1.1

$$(2.5) \quad Li_2(x) + Li_2(-u^2) = -\frac{1}{2} \log^2(1-x)$$

$$(2.6) \quad Li_2(u^2) + Li_2(-u^2) = \frac{1}{2} Li_2(u^4)$$

$$(2.7) \quad Li_2(u^4) = 2Li_2(u^2) - 2Li_2(x) - \log^2(1-x)$$

Adding (2.4) and (2.7) gives

$$(2.8) \quad Li_2(u^4) - Li_2(u^2) = Li_2(\bar{z}) - Li_2(z) - 3Li_2(x) + \frac{\pi^2}{3} - Li_2\left(\frac{1}{2}\right) + \log(2) \log(1-x) - \log^2(1-x) - \log(z) \log(1-z) - \frac{1}{2} \log(1+z) \log\left(\frac{1+z}{z^2}\right)$$

We want

$$(2.9) \quad \text{Im}(Li_2(u^4) - Li_2(u^2)) = \frac{1}{i}(Li_2(\bar{z}) - Li_2(z)) - \frac{3}{2i}(Li_2(x) - Li_2(\bar{x})) + \text{Im} \left\{ \log(2) \log(1-x) - \log^2(1-x) - \log(z) \log(1-z) - \frac{1}{2} \log(1+z) \log\left(\frac{1+z}{z^2}\right) \right\}$$

We get

$$\frac{1}{i}(Li_2(\bar{z}) - Li_2(z)) = \frac{1}{i} \sum_{n=1}^{\infty} \frac{1}{n^2} \left\{ \left(\frac{i}{\sqrt{8}}\right)^n - \left(\frac{-i}{\sqrt{8}}\right)^n \right\} = \frac{1}{\sqrt{2}} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} \left(-\frac{1}{2}\right)^{3n}$$

and using Lemma 2

$$\begin{aligned} & \frac{1}{2i}(Li_2(x) - Li_2(\bar{x})) = \frac{1}{2i}(Li_2\left(\frac{1-z}{2}\right) - Li_2\left(\frac{1+z}{2}\right)) \\ &= -\frac{z}{i} \sum_{n=1}^{\infty} \frac{H_n}{2n+1} z^{2n} + \frac{1}{4i} \left\{ Li_2\left(\frac{1-z}{2}\right) - Li_2\left(\frac{1+z}{2}\right) \right\} + \frac{\log(2)}{2i} \log\left(\frac{1-z}{1+z}\right) \end{aligned}$$

$$= \frac{1}{\sqrt{8}} \sum_{n=1}^{\infty} \frac{H_n}{2n+1} \left(-\frac{1}{2}\right)^{3n} + \alpha \log\left(\sqrt{\frac{9}{8}}\right) + \frac{\log(2)}{\sqrt{8}} \sum_{n=0}^{\infty} \frac{1}{2n+1} \left(-\frac{1}{2}\right)^{3n}$$

Putting everything together we obtain Broadhurst's formula (all logarithms cancel).

References.

1. D.Broadhurst, A dilogarithmic 3-dimensional tetrahedron, hep-th/980525
2. L.Lewin, Dilogarithms and associated functions, Macdonald, London 1958
3. A.P.Prudnikov, O.I.Marichev, Yu.A.Brychkov, Integrals and series I (Russian) Moskva Nauka, 1986

Institute of Algebraic Meditation
 Fogdaröd 208
 S-2433 Höör
 Sweden
 gert@maths.lth.se